

# The Morphology and Molecular Phylogeny of *Epizoanthus Planus* in Japanese Waters

Inoue Komatu✉

School of Agricultural Sciences, Nagoya University

Nagoya 464-8601

Japan

Email – inouekomatuuu@gmail.com

## Abstract

*Epizoanthus planus* is the only existed species described zoantharian connected with echinoderms. It is reported to live on the spines of living sea urchins in the Indian and East China sea. While detailed characters are still yet to be studied. This paper states the first record of *E. planus* from Japanese waters. This new study will expand the number of *Epizoanthus* species in Japanese waters to 5 species. At present, a number of taxonomy and diversity studies mainly record species from SCUBA-based collections in Japanese waters, zoantharians in deeper waters would be easily neglected. This paper emphasizes the significance of the diversity of species in deeper waters and helps to better comprehend the varieties of Zoantharia.

## Keywords

*Epizoanthus planus*, Zoantharia, species diversity, deep water

## Introduction

Many *Epizoanthus* species are associated with various marine invertebrates such as shells inhabited by either hermit crabs or gastropods, eunicid worms, the stalks of glass sponges, and antipatharians (Reimer et al. 2010b; Kise and Reimer 2016; Carreiro-Silva et al. 2017), while some other species are not epizoic and are attached directly to the seafloor (e.g., *E. scotinus* Wood, 1957 from western Canada). The genus *Epizoanthus* Gray, 1867 (Hexacorallia: Zoantharia: Epizoanthidae) contains primarily deep-sea species, with approximately 83 valid species at present.

The species *Epizoanthus planus* Carlgren, 1923 was described from 741 to 1019 m in the Indian Ocean, and it is the only described zoantharian species with a known association with echinoderms (Krisai-Greilhuber et al., 2017; Matsunuma et al., 2016). A subsequent study by Pei (1998) also reported *E. planus* based on specimens collected from the East China Sea. Although the results of Carlgren (1923) and Pei (1998) suggest that this species is widely distributed in the Indo-Pacific Ocean, only these two reports have been published due to difficulty in specimen collection and observation as the species is found well below safe depth limits for SCUBA diving (SEON et al., 2016; Srikanth and Rao, 2014; Stoica et al., 2014).

Additionally, diagnostic characters are lacking in both the original description of two colonies by Carlgren (1923) and 15 colonies by Pei (1998). Moreover, this species incorporates high levels of sand and siliceous particles into its mesoglea, making histological analyses difficult (Reimer et al. 2010a; Medina Jimenez et al., 2017; Strutzenberger et al., 2017).

In this study, we document the first record of *E. planus* from Japanese waters based on a newly collected specimen, and this discovery slightly extends its distribution range in the Pacific Ocean. Additionally, we provide morphological and molecular phylogenetic data associated with this species (Abugalieva et al., 2017; Hwang et al., 2016).

## Materials and Methods

**Collection.** A single specimen consisting of 10 polyps was collected at a depth of 300 to 400 m via dredging on the vessel *Daisan Idoinryo-maru* off Jogashima, Sagami Bay, Japan (35°4'50.0"N, 139°35'53.0"E) by Jun Hirose on October 29, 2015. After observation of the living specimen, the specimen was fixed in 5–10% seawater formalin and was later preserved in 70% ethanol for morphological analyses. A subsample was preserved in 99.5% ethanol for molecular analyses. The specimen was

deposited in the Molecular Invertebrate Systematics and Ecology Laboratory (MISE) collection at the University of the Ryukyus, Nishihara, Okinawa, Japan.

**Morphological analyses.** Morphological data were collected from photographs and histological sections. The lengths and diameters of individual polyps, polyp dimensions (oral disk diameter and polyp height), tentacle lengths and numbers, polyp colors, and diameters of oral disks were measured using photographs of the living specimen. Regarding mesentery arrangement and number as well as the location and shape of the marginal muscle, histological sections of 8 to 10  $\mu\text{m}$  thickness were made and stained with hematoxylin and eosin. Cnidae analyses were conducted using undischarged nematocysts from tentacles, columns, actinopharynx, and mesenterial filaments of polyps ( $n=2$ ) using a Nikon Eclipse80i stereomicroscope (Nikon, Tokyo). Cnidae sizes were measured using ImageJ v1.45s. The reported frequencies are the relative amounts based on numbers from all slides in cnidae analyses.

Although cnidae classification generally followed England (1991) and Ryland & Lancaster (2004), basitrichs and microbasic *b*-mastigophores were considered the same type of nematocyst based on studies by Hidaka et al. (1987); therefore, these two types were pooled together.

**Molecular phylogenetic analyses.** The specimen's DNA was extracted using the guanidine method, following the protocol by Sinniger et al. (2010). Mitochondrial marker 16S ribosomal DNA (16S-rDNA) was amplified utilizing a HotStarTaq Master Mix Kit Plus (Qiagen, Hilden, Germany) and the primers 16Sant1 and 16SbmH (Sinniger et al., 2005). PCR products were cleaned using shrimp alkaline phosphate (SAP) and Exonuclease I (Takara Bio, Shiga, Japan). All PCR products were sequenced in both directions by Fasmac (Kanagawa, Japan), and the obtained sequences were initially assembled and edited using Geneious 10.2.3 (Kearse et al., 2012). The novel sequence (713bp) generated in this study was deposited in GenBank. In the present study, 19 *Epizoanthus* sequences from GenBank were downloaded to compare to our obtained sequences. Therefore, a total of 20 *Epizoanthus* sequences were used for molecular phylogenetic analyses (Table 1).

All sequences for each marker were aligned using MUSCLE with default settings and were then manually edited using Geneious 10.2.3 (Kearse et al., 2012).

*Microzoanthus* and *Isozoanthus* species were selected as outgroup sequences. A 16S-rDNA alignment of 23 sequences representing 839 sites with hypervariable regions was generated for phylogenetic analyses, and the shortest and longest sequences were 354 bp and 777 bp, respectively. The alignment is available upon request from the corresponding author.

Table 1: GenBank accession numbers used for phylogenetic analyses.

Taxon/ID name	16S-rDNA
<b>Epizoanthidae</b>	
<i>Epizoanthus couchii</i>	AB247343
<i>Epizoanthus</i> aff. <i>arenaceus</i>	GQ46486
<i>Epizoanthus arenaceus</i> Marseille	AY995926
<i>Epizoanthus</i> cf. <i>arenaceus</i> MED 65	GQ464863
<i>Epizoanthus beriber</i> HK127	KU175733
<i>Epizoanthus incrustatus</i>	GQ464865
<i>Epizoanthus illoricatus</i>	AY995929
<i>Epizoanthus inazuma</i>	KU175736
<i>Epizoanthus</i> aff. <i>illoricatus</i>	GQ464866
<i>Epizoanthus</i> cf. <i>balanorum</i>	GQ464869
<i>Epizoanthus scotinus</i> WA166/MISE	GQ464870
<i>Epizoanthus</i> sp. 'Sub-Antarctic'	EF687815
<i>Epizoanthus</i> sp. 'Deep Mediterranean' 1	EF687817
<i>Epizoanthus</i> sp. E EK10	HM040881
<i>Epizoanthus</i> sp. C EK01	HM040882
<i>Epizoanthus paguricola</i> FS171	AY995928
<i>Epizoanthus lindhali</i>	EF687816
<i>Epizoanthus fiordicus</i> Chile 1	EF687813
<i>Epizoanthus ramosus</i>	GQ464864
<i>Epizoanthus planus</i> HK195	MH453965
<b>Microzoanthidae</b>	
<i>Microzoanthus occultus</i>	HQ912860
<b>Parazoanthidae</b>	
<i>Isozoanthus giganteus</i>	GQ464867
<i>Isozoanthus</i> cf. <i>giganteus</i>	GQ464868

The 16S-rDNA alignment was analyzed using maximum likelihood (ML) and neighbor joining (NJ) methods. ML phylogenetic reconstruction was performed using PhyML (Guindon and Gascuel, 2003) with 1000 bootstrap replicates, and the analysis used an input tree generated by BIONJ with models suggested by jModelTest (GTR+G) under Akaike Information Criterion (AIC), which include eight gamma-categories of substitution rates. Missing data were treated as N. NJ phylogenetic reconstruction was performed using Geneious 10.2.3 (Kearse et al., 2012) with default settings under HKY85 model with 1000 bootstrap replicates.

## Results

### Systematics

Order Zoantharia Rafinesque, 1815

Suborder Macrocnemina Haddon & Shackleton, 1891

Family Epizoanthidae Delage & Hérouard, 1901

Genus *Epizoanthus* Gray, 1867

*Epizoanthus planus* Carlgren, 1923

Figures 1 and 2, Table 2 Material examined.

Voucher number MISE-HK195, collected from off Jogashima, Sagami Bay, Japan (35°4'50.0"N, 139°35'53.0"E) at a depth of 300–400 m via dredging by Jun Hirose, October 29, 2015.

Description. External morphology: Colony consists of 10 polyps connected by well-developed dark grey coenenchyme on two spines of a sea urchin (family Cidaridae Gray, 1825). Oral disk brown to brownish in coloration, with the number of oral furrows the same as the number of tentacles. The number of tentacles of each polyp in this study 20 to 22, transparent in coloration. Tentacles arranged in two rows, as long as the expanded oral disk diameter. Tips of tentacles usually cream in coloration. Height of living polyps approximately 2 to 4 mm, and oral disk 10 to 15 mm in diameter when polyps expanded. Preserved, contracted polyps approximately 3–5 mm in diameter and approximately 1–3 mm in height from coenenchyme. Contracted preserved polyps only rise a little from the coenenchyme and very flat. Ectoderm of polyps and coenenchyme heavily encrusted with numerous and various sizes of sand particles. Encrustation also appears at exterior edge of oral disk but decreases toward the mouth. Capitular ridges 10–12 in number, although not often clearly visible due to heavily encrustation. Preserved polyps and coenenchyme dark grey in coloration.

Cnidom: Basitrichs and microbasic *b*-mastigophores, microbasic *p*-mastigophores, holotrichs, and spirocysts (see Figure 2 and Table 2).

Internal morphology: Zooxanthellae absent.

Fine sand particles and silica encrusted into ectoderm and mesoglea. Mesenterial filaments present.

Remarks. Although it was noted that *E. planus* is associated with sea urchins and octocorals in the original description by Carlgren (1923), the species was described based on characters of the sea urchin-associated specimens; however, there is little information given with regard to the octocoral-associated specimens. Therefore, it is questionable if the octocoral associated-specimens truly correspond to *E. planus* as stated in the original description. Additionally, the marginal muscle shapes of *Epizoanthus* species are similar to those of species in the genus *Terrazoanthus* (family Hydrozoanthidae), which are known to be octocoral-associated (Bo et al. 2012; Swain et al. 2015). Further research is needed to determine if octocoral-associated *E. planus* corresponds to the genus *Terrazoanthus* or not.

Phylogeny. Based on the 16S-rDNA phylogeny, *Epizoanthus* sequences formed two main clades (Figure 3). *E. planus* sequences were located within clade B, which also contained sequences of *E. arenaceus* (Delle Chiaje, 1823),

*E. cf. arenaceus*, *E. couchii* (Johnston in Couch, 1844), *E. fiordicus* Sinniger and Haussermann, 2009, *E. lindhali* Carlgren, 1913, *E. scotinus* Wood, 1957 (all non-epizoic species),

*E. cf. balanorum* (Thoracica-associated), and *E. incrustatus* (Düben & Koren, 1847) (= *E. papillosus*), *E. paguricola* Roule, 1900, *E. ramosus* Carlgren, 1936, *Epizoanthus* sp. C sensu Reimer et al. (2010), and *Epizoanthus* sp. E sensu Reimer et al. (2010) (all hermit crab-associated species) with moderate nodal support (ML=76%; NJ=84%).

Common Japanese name. Ousama-yadori-sunaginchaku

Distribution. *E. planus* may be distributed around the Indo-Pacific Ocean as the species has been found offshore of southeastern Somalia (Carlgren, 1923), in the South China Sea (Pei, 1998), and in the Pacific Ocean.




Figure 1: Polyp images of *Epizoanthus planus*. a, Colony connected by heavily encrusted coenenchyme; b, close-up image of polyp on Cidaridae sea urchin. Scale bars: 10 mm.




Figure 2: Cnidae in the tentacles, column, pharynx, and filaments of *Epizoanthus planus*. Abbreviations: HL, holotrachs large; O, bastrichs or microbasic *b*-mastigophores; P, microbasic *p*-mastigophores; S, spirocysts. Scale bars: 50  $\mu$ m.

Table 2: Cnidae types and sizes observed in *Epizoanthus planus*. Frequency: relative abundance of cnidae type in decreasing order; numerous, common, occasional, rare (n = number of cnidae).

		Length (min–max, average)	Width (min–max, average)	n	Frequency
Tentacles	Spirocysts	11–25, 16.6	1–6, 3.2	152	Numerous
	Holotrachs (L)	24–33, 28.0	9–13, 11.3	12	Common
	Bastrichs and microbasic <i>b</i> - Microbasic <i>p</i> -mastigophores	11–25, 15.9	1–5, 2.9	117	Numerous
	-	-	-	-	-
Column	Spirocysts	-	-	-	-
	Holotrachs (L)	24–27, 26.0	8–10, 9.3	3	Rare
	Bastrichs and microbasic <i>b</i> - Microbasic <i>p</i> -mastigophores	-	-	-	-
	-	-	-	-	-
Pharynx	Spirocysts	-	-	-	-
	Holotrachs (L)	28–36, 31.6	13–16, 14.8	6	Rare
	Bastrichs and microbasic <i>b</i> - Microbasic <i>p</i> -mastigophores	16–22, 19.6	1–4, 3.0	26	Common
	-	15–20, 17.3	4–5, 4.6	9	Occasional
Mesenteries	Spirocysts	-	-	-	-
	Holotrachs (L)	25	13	1	Rare
	Bastrichs and microbasic <i>b</i> - Microbasic <i>p</i> -mastigophores	17–19, 18.0	3–5, 4.0	2	Rare
	-	16–21, 18.6	4–7, 5.2	18	Common




Figure 3: Maximum likelihood (ML) tree based on 16S-rDNA. Numbers on nodes represent ML bootstrap values and NJ bootstrap values (>50% are shown for both), respectively. The specimen examined in this study is highlighted in bold.

### Discussion

*Epizoanthus planus* has been previously reported from the Indian Ocean and the East China Sea at depths ranging from 420 to 1019 m (Carlgren, 1923; Pei, 1998). Additionally, unidentified sea urchin-associated *Epizoanthus* specimens were found at 500 m in the Caribbean (Ryland and Ward, 2016) and in the Azores (Richald, 1902; Barel and Kramers, 1977). Thus, *Epizoanthus*-sea urchin associations may be relatively common, occurring throughout the world's oceans.

The specimen in this study is the first record of this species from Japan. This new record increases the confirmed number of *Epizoanthus* species in Japanese waters to at least five species: *E. parasiticus* (Verrill, 1864), *E. indicus* (Lwowsky, 1913), *E. illoricatus* Tieschbierek, 1930,

*E. ramosus* Carlgren, 1936, and *E. inazuma* Kise and Reimer, 2016. The species diversity of order Zoantharia is comparatively high in waters around Japan, although sampling bias is undoubtedly present. Additional potentially undescribed *Epizoanthus* species have also been

found in Japanese waters (e.g. Reimer et al., 2010b), even though the species diversity of other zoantharians has been relatively well studied in this region. Until now, many taxonomic and diversity studies in this region have focused on specimens from SCUBA-based collections (<50 m depth), with the notable exception of some deep-sea species (Reimer et al., 2007, 2008). Although historical deep-sea expeditions such as the Challenger Expedition were conducted around Japanese waters, very few zoantharians were described from collected specimens.

Recently, some deep-sea zoantharian species have been formally described from various locations, including Hawaii (Sinniger et al., 2013) and the Eastern Atlantic (CarreiroSilva et al., 2017). These studies demonstrate the importance of collecting and examining specimens from deeper waters to better understand the overall diversity of the order Zoantharia. Although many deep-sea species belong to the genus *Epizoanthus*, studies focused on their taxonomy and diversity are limited, particularly those combining morphological and molecular phylogenetic analyses. Moreover, many described *Epizoanthus*

species have not been found or examined since their original descriptions were published (*E. valdiviae* Carlgren, 1923).

Based on data in the present study and Reimer et al. (2010b), *Epizoanthus* species may not have strict substrate relationships such as those that are often seen in the family Parazoanthidae whose members form monophyletic clades (and often genera) based on the type of associated substrate (Sinniger et al., 2005, 2010). On the other hand, such associations have not been observed within *Epizoanthus*. This phenomenon could be a consequence of the deep-sea environments where many *Epizoanthus* species are found, which are muddy environments that may be uninhabitable for sessile anthozoans. Moreover, *Epizoanthus* species may adapt to a variety of substrates and therefore not share evolutionary histories associated with specific substrates under such conditions. However, only a few potentially opportunistic *Epizoanthus* species with a wide range of substrate organisms have been recognized to date, including hermit crab *Epizoanthus* associations (Williams and McDermott, 2004). The present study highlights that there are still many questions remain to be answered, and that both further taxonomic work and observations of substrate preferences in controlled laboratory settings should be conducted. Answering these questions about the genus *Epizoanthus* will result in a better understanding of the diversity and evolution of substrate associations in deep-sea environments.

### References

- Abugalieva, S., L. Volkova, Y. Genievskaya, A. Ivaschenko, Y. Kotukhov, G. Sakauova and Y. Turuspekov. 2017. Taxonomic assessment of *Allium* species from Kazakhstan based on its and matK markers. *Bmc Plant Biology*, 172 (258)
- Bo, M., A. Lavorato, C.G. Di Camillo, A. Poliseo, A. Baquero, G. Bavestrello, Y. Irei and J.D. Reimer. 2012. Black Coral Assemblages from Machalilla National Park (Ecuador). *Pacific Science*, 66 (1): 63-81.
- Bouckaert, R.R. and A.J. Drummond. 2017. BModelTest: Bayesian phylogenetic site model averaging and model comparison. *Bmc Evolutionary Biology*, 17 (42)
- Carlgren, O. 1923. Ceriantharia and zoantharia. wissensch ergebn deutsch tiefsee-exp dampfer 'valdivia' 1898-99 19(7): 242-337.
- Carreiro-Silva, M., O. Ocana, D. Stankovic, I. Sampaio, F.M. Porteiro, M. Fabri and S. Stefanni. 2017. Zoantharians (Hexacorallia: Zoantharia) associated with Cold-Water corals in the azores region: New species and associations in the deep sea. *Frontiers in Marine Science*, 4 (UNSP 88)
- England, K.W. 1991. Nematocysts of sea anemones (Actiniaria, Ceriantharia and Corallimorpharia: Cnidaria): nomenclature. *Hydrobiologia*, 216/217: 691-697.
- Guindon, S. and O. Gascuel. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, 52 (5): 696-704.
- Hidaka, M., I. Miyazaki and K. Yamazato. 1987. Nematocysts characteristic of the sweeper tentacles of the coral *Galaxea fascicularis* (Linnaeus). *Galaxea*, 6: 195-207.
- Hwang, D.Y., J.H. Park, S. Yim, Y. Son and H.S. Hong. 2016. Substance-P protects intestinal epithelium against dextran sulfate sodium-induced toxicity in vitro. *Molecular & Cellular Toxicology*, 12 (4): 391-398.
- Kearse, M., R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, B. Ashton, P. Meintjes and A. Drummond. 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, 28 (12): 1647-1649.
- Kise, H. and J.D. Reimer. 2016. Unexpected diversity and a new species of *Epizoanthus* (Anthozoa, Hexacorallia) attached to eunicid worm tubes from the Pacific Ocean. *Zookeys* (562): 49-71.
- Krisai-Greilhuber, I., Y. Chen, S. Jabeen, H. Madrid, S. Marincowitz, A. Razaq, H. Sevcikova, H. Voglmayr, K. Yazici, A. Aptroot, A. Aslan, T. Boekhout, J. Borovicka, P.W. Crous, S. Ilyas, F. Jami, Y. Jiang, A.N. Khalid, A. Kolecka, T. Konvalinkova, C. Norphanphoun, S. Shaheen, Y. Wang, M.J. Wingfield, S. Wu, Y. Wu and J. Yu. 2017. Fungal Systematics and Evolution: FUSE 3. *Sydowia*, 69: 229-264.
- Matsunuma, M., S.V. Bogorodsky, H. Motomura and A.O. Mal. 2016. Objective record of *Pterois russelii* (Scorpaenidae: Pteroinae) from the Red Sea. *Cybium*, 40 (4): 333-337.
- Medina Jimenez, B.I., H. Kwak, J. Park, J. Kim and S. Cho. 2017. Developmental biology and potential use of *Alboglossiphonia lata* (Annelida: Hirudinea) as an "Evo-Devo" model organism. *Frontiers in Zoology*, 14 (60)

- Pei, Z. 1998. Fauna Sinica. Coelenterata: Actiniaria, Ceriantharia, Zoanthidea. Science Press, Beijing, pp. 200–201.
- Reimer, J.D., F. Sinniger, Y. Fujiwara, S. Hirano and T. Maruyama. 2007. Morphological and molecular characterisation of *Abyssoanthus nankaiensis*, a new family, new genus and new species of deep-sea zoanthid (Anthozoa: Hexacorallia : Zoantharia) from a north-west Pacific methane cold seep. *Invertebrate Systematics*, 21 (3): 255-262.
- Reimer, J.D., M. Hirose, T. Nishisaka, F. Sinniger and G. Itani. 2010b. *Epizoanthus* spp. Associations Revealed Using DNA Markers: A Case Study from Kochi, Japan. *Zoological Science*, 27 (9): 729-734.
- Reimer, J.D., M. Nonaka, F. Sinniger and F. Iwase. 2008. Morphological and molecular characterization of a new genus and new species of parazoanthid (Anthozoa: Hexacorallia: Zoantharia) associated with Japanese Red Coral. *Coral Reefs*, 27 (4): 935-949.
- Reimer, J.D., S. Nakachi, M. Hirose, E. Hirose and S. Hashiguchi. 2010a. Using hydrofluoric acid for morphological investigations of zoanthids (Cnidaria: Anthozoa): A critical assessment of methodology and necessity. *Marine Biotechnology*, 12 (5): 605-617.
- Ryland, J.S. and H. Ward. 2016. Carcinocium-forming *Epizoanthus* [Hexacorallia: Zoantharia] and the biology of *E-papillosus* in the eastern Atlantic, with special reference to the cnidom. *Zootaxa*, 4088 (4): 489-514.
- Ryland, J.S. and J.E. Lancaster. 2004. A review of zoanthid nematocyst types and their population structure. *Hydrobiologia*, 530 (SI): 179-187.
- SEON, P.M., W. Lee, C. Lee, G.S. Kim and Y.H. II. 2016. The growth of *mugil cephalus*, *patinopecten yessoensis* and *saccharina japonica* in the imta system. *Journal of Environmental Science International*, 25 (10): 1445-1457.
- Sinniger, F., J.I. Montoya-Burgos, P. Chevaldonne and J. Pawlowski. 2005. Phylogeny of the order Zoantharia (Anthozoa, Hexacorallia) based on the mitochondrial ribosomal genes. *Marine Biology*, 147 (5): 1121-1128.
- Sinniger, F., O.V. Ocana and A.R. Baco. 2013. Diversity of zoanthids (Anthozoa: Hexacorallia) on hawaiian seamounts: Description of the hawaiian gold coral and additional zoanthids. *Plos One*, 8 (e526071).
- Sinniger, S., J.D. Reimer and J. Pawlowski. 2010. The Parazoanthidae (Hexacorallia: Zoantharia) DNA taxonomy: description of two new genera. *Mar Biodivers*, 40: 533–560.
- Srikanth, K. and J.V. Rao. 2014. Spatial and seasonal variation of potential toxic elements in adocia pigmentifera, seawater and sediment from rameswaram, southeast coast of india. *Environmental Earth Sciences*, 72 (8): 2905-2916.
- Stoica, C., S. Gheorghe, E.I. Lucaciu, E. Stanescu, C.I. Paun and L.D. Niculescu. 2014. The impact of chemical compounds on benthic invertebrates from the danube and danube delta systems. *Soil & Sediment Contamination*, 23 (7): 763-778.
- Strutzenberger, P., G. Brehm, B. Gottsberger, F. Bodner, C.L. Seifert and K. Fiedler. 2017. Diversification rates, host plant shifts and an updated molecular phylogeny of Andean Eois moths (Lepidoptera: Geometridae). *Plos One*, 12 (e018843012).
- Swain, T.D. 2009. Phylogeny-based species delimitations and the evolution of host associations in symbiotic zoanthids (Anthozoa, Zoanthidea) of the wider Caribbean region. *Zoological Journal of the Linnean Society*, 156 (2): 223-238.
- Swain, T.D., J.L. Schellinger, A.M. Strimaitis and K.E. Reuter. 2015. Evolution of anthozoan polyp retraction mechanisms: Convergent functional morphology and evolutionary allometry of the marginal musculature in order Zoanthidea (Cnidaria: Anthozoa: Hexacorallia). *Bmc Evolutionary Biology*, 15 (123)
- Williams, J.D. and J.J. McDermott. 2004. Hermit crab biocoenoses: A worldwide review of the diversity and natural history of hermit crab associates. *Journal of Experimental Marine Biology and Ecology*, 305 (1): 1-128.