

Effects of Chitosan Addition on Weaned Piglets

Emma Bland
 Faculty of Science, Murdoch University
 Murdoch 6150
 Australia
 Email - Emmbla@au.dk

Abstract

The purpose of this experiment was to determine the commercial sources of chitosan (CS) in weaned pigs to enhance performance, antioxidant function and immune response. On 28 ± 2 days, 60 crossbred piglets (Duroc \times Landrace \times Yorkshire) with an average live weight of 8.85 ± 1.52 kg were weaned and randomly divided into 5 treatment groups, which were fed with 0 (basic diet, control) and 250, 500, 1000, and 2000 mg/kg CS. The experiment lasted two weeks. Daily food intake was recorded and calculated. The consistency of feces throughout the period was monitored. Two weeks later, blood samples were collected and the antioxidant and immune parameters were determined. The results showed that CS increased average daily gain and daily gain: daily feed intake during the experiment. CS improved the average fecal score in the second week. Compared with the control group, the average score decreased. CS can increase the total antioxidant capacity in serum, superoxide dismutase, catalase and glutathione peroxidase activity as well as reduced glutathione content, and reduce serum malondialdehyde and cortisol contents. In addition, CS increased serum IL-1 β , IL-2 and IgG levels. These findings indicate that the use of CS improves performance and antioxidant function and regulates the immune response of weaned pigs.

Keywords

Chitosan; Anti-oxidative capability; Immune response; Piglets.

Introduction

Weaning, during which piglets are subjected to abrupt changes in environment, behaviour and diet, is one of the most stressful events in a pig's life (Lee et al., 2012). It can contribute to gastrointestinal disorders, immune system dysfunctions, and endocrine system imbalance, and may cause diarrhoea, increased susceptibility to disease, morbidity, and mortality, and inhibited growth and health (Zhang et al., 2015). Weaning therefore causes serious economic losses in pig production worldwide (Hena-Diaz et al., 2018). Additionally, a recent study demonstrated that weaning could induce oxidative stress, resulting in oxidative damage in pigs (Liu et al., 2017).

Reducing post-weaning stress is one of the main challenges for the pig industry (Siel et al., 2016). In traditional commercial practice, the use of additives has been recommended as a way to ameliorate or relieve post-weaning stress during this stage (Liu et al., 2016). Among them, CS has been indicated as having positive effects on pig growth and health. CS is obtained by deacetylation of chitin, which is present in exoskeletons of arthropods such as crustaceans and insects, and in the outer cell wall of fungi (Xiong et al., 2014). Compared with other natural polysaccharides, a plethora of applications of this biopolymer have been found owing to its high biocompatible, biodegradable, and nontoxic

nature (Inaba et al., 2013). As a natural cationic polysaccharide, CS contains reactive functional groups, that is, hydroxyl and amino groups. Numerous studies have demonstrated that CS has unique biological activities, which include anti-microbial, immunostimulatory, anti-inflammatory, anti-tumour, anti-hypertension, hypocholesterolemic, anti-obesity, and anti-oxidation properties (Calinescu et al., 2012). Notably, because of these properties, CS can be used as a pro-growth and health feed supplement for farm animals. However, the causal link between CS and the improvement in pig health status remains largely unknown. Hence, further clarification is necessary (Lv et al., 2011).

Accordingly, the main objective of this work was to investigate the effects of CS on anti-oxidative function and immune response in weaned pigs (Han et al., 2011). These findings may be useful in elucidating mechanisms that promote growth and health through enhanced anti-oxidative and immune functions in animals that receive dietary supplementation with CS.

Materials and Methods

The experiment was carried out in the Experimental Farms of Inner Mongolia Agricultural University in China and received prior approval from the Institutional Animal Care and Use Committee of this Institution.

The experiment followed a randomized block design. A total of 60 healthy weaned pigs (Duroc × Yorkshire × Landrace, 30 barrows and 30 gilts) from 12 litters (five pigs per litter) were selected from a commercial herd. The animals were weaned at the age of 28 days with average initial bodyweight of 8.85 (± 1.52) kg. The animals were randomly assigned to five treatments ($n=12$, six barrows and six gilts), and were housed in 60 pens (one pig per pen) taking initial body weight and litter (one pig from each litter per treatment)

into account. The dietary treatments consisted of a basal diet (control treatment) that was formulated to meet the nutrient requirements of the National Research Council. The compositions are shown in Table 1. Treatments 2-5 consisted of the basal diet supplemented with 250, 500, 1000 and 2000 mg CS/kg. No medication or other additives were included in the diets. Throughout the experimental period, animals were allowed access to feed and water ad libitum and exposed to natural lighting.

Table 1: Composition and chemical analysis of basal experimental diets (air dry basis)

Ingredients	Content	Nutrient levels	Content
Soybean meal	26.57	Crude protein (N×6.25)	20.37
Wheat bran	5.00	Lysine	1.45
Fish meal	3.00	Methionine	0.47
Soybean oil	1.00	Methionine + cysteine	0.87
Limestone	1.39	Threonine	0.96
Dicalcium phosphate	0.89	Tryptophan	0.24
Salt	0.30	Calcium	0.90
L-Lysine HCL	0.45	Phosphorus	0.61
DL-Methionine	0.16		
L-Threonine	0.15		
Vitamin/mineral premix 1	1.00		
Total	100.00		

The pigs were weighed at the beginning (day 1) and end (day 14) of the feeding period to obtain bodyweight and to calculate average daily bodyweight change.

The faecal consistency was classified using a score ranking from 1 to 5 described by Pierce *et al.* as 1) hard firm faeces; 2) slightly soft faeces; 3) soft, partially formed faeces; 4) loose, semi-liquid faeces (diarrhoea); and 5) watery mucous-like faeces (serious diarrhoea). Faeces scores were given daily for individual pigs from day 1 until day 14. These scores were collected by a trained individual with no prior knowledge of the dietary treatment.

At the end of the experiment, after an overnight fast, six animals (three barrows and three gilts) were selected from each treatment at random and blood samples (10 mL per pig) were collected via anterior vena cava into non-heparinized vacuum tubes. These blood samples were centrifuged (1200 × g, 4°C) for 10 min to separate serum. Serum was stored at -20 °C until further assays.

In the present study, serum cortisol level was determined with commercial ELISA kits (Nanjing Jiancheng Bioengineering Institute, Nanjing, China) according to the manufacturer's instructions. All samples were analysed in duplicate.

The effects of CS ingestion on the antioxidant capacities in the present study were determined by measuring antioxidant-related indices in the serum, such as the contents of total anti-oxidative capability (T-AOC), enzymatic activity of catalase (CAT), total superoxide dismutase (T-SOD), copper/zinc superoxide dismutase (CuZn-SOD), glutathione peroxidase (GPx), and reduced glutathione (GSH) and malondialdehyde (MDA), using commercial assay kits according to the manufacturers' instructions (Nanjing Jiancheng

Bioengineering Institute, Nanjing, China). All assays were carried out in duplicate with samples from six pigs per group. Determination of T-AOC enabled the total activities of several parameters to be evaluated, including vitamins (e.g. vitamin C and vitamin E), amino acids (e.g. cysteine, methionine, tryptophan, and histidine) and metal protein (e.g. transferrin and lactoferrin), β -carotene, and GSH, all of which can convert Fe^{3+} to Fe^{2+} . Then Fe^{2+} can be combined with phenanthroline to form stable coloured chelates. The T-AOC was measured with a spectrophotometer at 520 nm and expressed as activity unit U/mL serum. In this assay, 1 U T-AOC was defined as a 0.01 increase in the absorbance value at 37°C for 1 min per mL serum. In the present assay, superoxide anion radicals (O_2^-) are generated by the xanthine/xanthine oxidase system, which reacts with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyltetrazolium chloride to form a purple formazan dye, which is used as the detector. The SOD activity was measured at 550 nm with a spectrophotometer and 1 U of T-SOD activity was defined as the amount of serum capable of inhibiting the reduction of nitro blue tetrazolium by 50% of maximum inhibition. The CuZn-SOD activity was measured by adding sodium cyanide (NaCN) to the reaction system to inactivate manganese superoxide dismutase (Mn-SOD) activity, while Mn-SOD was calculated by subtracting the CuZn-SOD value from T-SOD. The CAT activity was determined as the decrease in hydrogen peroxide. The enzymatic reaction of H_2O_2 degraded by CAT was terminated by the addition of ammonium molybdate, and the rest H_2O_2 combined with ammonium molybdate, generated a light-yellow complex that could be measured spectrophotometrically at 405 nm. The CAT activity was expressed as U/mL serum, and 1U of CAT activity was defined as the amount of enzyme needed to

decrease 1 μmol of H_2O_2 at 37°C for 1 sec per mL serum. The GPx activity was detected by quantifying the reduction rate of reduced glutathione. The GSH reacts with 5, 5-dithiobis nitrobenzoic acid to form yellow compounds. The GPx was determined with a spectrophotometer at 412 nm and GPx activity was expressed as U/mL serum, in which 1U of GPx was expressed as the amount that reduced the level of GSH by 1 $\mu\text{mol/L}$ in 5 min per 0.1 mL serum at 37°C . GSH can react with 5, 5'-dithiobis-2-nitrobenzoic acid to produce oxidized glutathione (GSSG) and yellow chromophore 2-nitro-5-thiobenzoic acid (TNB), which can be detected at 412 nm. GSH content was expressed as mg/L serum. Serum MDA content was quantified using the thiobarbituric acid method, whereby MDA reacts with thiobarbituric acid in an acidic medium at 95°C for 40 min to form a pink coloured product. MDA was determined with a spectrophotometer at 532 nm and MDA was expressed as nmol/mL serum.

The immune response status in serum was estimated by measuring the concentrations of cytokines (interleukin (IL)-1, IL-2, IL-4, IL-6, tumour necrosis factor (TNF)- α and γ -Interferon (IFN) and the immunoglobulins (IgA, IgG and IgM). These parameters were measured with the commercial porcine colorimetric ELISA kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, China) according to the manufacturer's instructions. All samples were analysed in duplicate.

The experimental data generated in this study were subjected to statistical analysis with the general linear model procedure of SAS (SAS Institute Inc., Cary, NC, USA). The one-way analysis of variance test was used

to analyse all parameters. In addition, data were analysed for linear and quadratic responses to increasing dietary CS levels (0, 250, 500, 1000 and 2000 mg CS/kg diet) using the GLM procedure of SAS. The individual pen represented the experimental unit. All results were presented in the tables as means \pm standard error of means (SEM). Probability values of $p < 0.05$ were used as the criterion for statistical significance, and $p < 0.10$ was considered a tendency.

Results

As shown in Table 2, there were linear or quadratic responses in daily gain ($p < 0.05$, $p < 0.01$) and G: F ratio ($p < 0.01$, $p < 0.01$) to the increasing amount of CS inclusion during the overall experimental period (days 0 to 14), with all the CS-supplemented treatments improving daily gain ($p < 0.01$) and G: F ratio ($p < 0.01$) compared with the control. However, the amount of CS in the diet had no effect ($p > 0.05$) on feed intake (ADFI) during the overall experimental period.

There were no differences in faecal score among the treatments from days 0 to 7 ($p > 0.05$). There was a quadratic decrease in faecal inconsistency as the level of CS was increased from day 8 to day 14 ($p < 0.05$) and during the overall period ($p < 0.05$) with all the CS-supplemented treatments having an improved faecal score from day 8 to day 14 ($p < 0.05$), and 500 and 1,000 mg/kg CS treatments improved faecal score during the overall experimental period compared with the control ($p < 0.05$).

Table 2: Effect of feeding chitosan on performance and faecal score of weaned piglets

Dietary treatment	Chitosan levels g/kg diet					SEM	Significance ³		
	0(control)	250	500	1,000	2,000		T	L	Q
Initial weight, kg	8.82	8.84	8.83	8.85	8.89	0.44	ns	ns	ns
Final weight, kg	12.62	13.61	14.11	13.90	13.64	0.54	ns	ns	ns
ADG, g/d	271.4 ^b	340.6 ^a	365.5 ^a	360.4 ^a	342.9 ^a	15.3	**	*	**
ADFI, g/d	576.5	587.6	613.2	596.8	586.7	19.3	ns	ns	ns
G:F g/g	0.47 ^b	0.58 ^a	0.60 ^a	0.60 ^a	0.58 ^a	0.03	**	**	**
	Faecal score (days) ²								
1-7	2.74	2.60	2.34	2.54	2.49	0.11	ns	ns	ns
8-14	2.66 ^a	2.20 ^b	2.09 ^b	2.03 ^b	2.20 ^b	0.12	*	ns	*
Overall (1-14)	2.70 ^a	2.40 ^{ab}	2.21 ^b	2.29 ^b	2.34 ^{ab}	0.10	*	ns	*

As shown in Figure 1, serum cortisol concentration was quadratic decreased ($p < 0.05$) as the level of dietary CS was increased, and all the CS-supplemented treatments had a decreased cortisol level ($p < 0.05$).

Values are means (six pigs per treatment) with standard deviations represented by vertical bars. Control: basal diet; CS-250: basal diet with 250 mg chitosan/kg; CS-500: basal diet with 500 mg chitosan/kg; CS-1000: basal diet with 1000 mg chitosan/kg; CS-2000: basal diet with 2000 mg chitosan/kg.




Figure 1: Effect of feeding chitosan on serum cortisol of weaned piglets.

As shown in Table 3, there was a quadratic increase response in activity of T-AOC ($p < 0.05$), CAT ($p < 0.05$), and content of GSH ($p < 0.05$) to increased CS inclusion, with all the CS-supplemented treatments having increased serum T-AOC ($p < 0.05$) and CAT ($p < 0.01$) activities, and GSH concentration ($p < 0.05$). Meanwhile, there were quadratic increase responses in SOD (T-SOD: $p < 0.01$, CuZn-SOD: $p < 0.10$, Mn-SOD: $p < 0.10$) to the increased CS, with all the CS-supplemented treatments except for 250 mg/kg

increasing serum T-SOD activity ($p < 0.05$) compared with the control. In addition, serum GPx activity was linearly ($p < 0.05$) or quadratically ($p < 0.01$) increased with increased CS, and all the CS-supplemented treatments except for 250 mg/kg increased GPx activity ($p < 0.05$) compared with the control. However, the concentration of serum MDA level was quadratically decreased ($p < 0.05$) with the increased amount of CS inclusion, and 500 and 2,000 mg/kg CS treatments had lower concentrations of MDA than the control.

Table 3: Effect of feeding chitosan on serum antioxidant enzymes reduced glutathione and malondialdehyde of weaned piglets

Dietary treatment	Chitosan levels g/kg diet					SEM	Significance ³		
	0 (control)	250	500	1,000	2,000		T	L	Q
T-AOC, U/mL	3.07 ^b	3.68 ^a	3.81 ^a	3.65 ^a	3.68 ^a	0.18	*	a	*
T-SOD	101.3 ^b	108.7 ^{ab}	120.2 ^a	117.3 ^a	114.3 ^a	4.3	*	a	**
CuZn-SOD	59.4	63.7	68.2	65.6	65.9	2.4	a	a	
Mn-SOD	41.8	45.0	52.0	51.7	48.4	3.5	a	a	
CAT, U/mL	2.66 ^b	3.79 ^a	3.79 ^a	3.48 ^a	3.50 ^a	0.21	**		a
GPx, U/mL	485.3 ^b	522.3 ^{ab}	548.0 ^a	548.0 ^a	543.2 ^a	16.3	*		*
GSH, mg/L	1.75 ^b	2.11 ^a	2.34 ^a	2.14 ^a	2.16 ^a	0.13	*	a	*
MDA, nmol/mL	3.64 ^a	3.24 ^{ab}	3.14 ^b	3.26 ^{ab}	3.17 ^b	0.14	*	a	*

Discussion

Undoubtedly, the negative effects of weaning stress on growth and health in piglets are known. Chitosan is a polysaccharide with non-nutritional properties, is not digested by mammalian enzymes, and is delivered to the large intestinal tract. However, numerous studies have demonstrated that CS might act as a growth promoter for farm animals. Yin *et al.* indicated that dietary CS (0.025%) fed to early weaned piglets increased their performance. In agreement with those findings, the present results showed that supplementation with CS significantly increased weight gain and gain/feed of weaned piglets. The authors' previous work verified an improvement in the small intestinal morphology and digestive enzyme activities such as amylase in CS-fed piglets, which could contribute to higher nutrient digestibility. On the other hand, a higher growth performance inversely supported improved nutrient digestibility. In this study, the faecal score decreased in pigs that were fed various amounts of CS compared with the control, which may also support the improvement in performance achieved during the experiment.

Cortisol is the main active hormone of the hypothalamic-pituitary-adrenal (HPA) axis in pigs. This is a cholesterol-derived steroid that is synthesized in the fascicular zone of the adrenal cortex under the control of the pituitary adrenocorticotropic hormone. Cortisol has proteolytic and lipolytic activity in the peripheral tissues and gluconeogenesis and protein synthesis activity in the liver. The HPA axis in pigs is exquisitely sensitive to endogenous and exogenous stimuli, both acute and chronic. It is therefore a valuable index for

monitoring stress conditions of farm animals. An elevation of cortisol illustrates the activation of HPA axis. The current experiment showed that the content of cortisol decreased significantly in piglets fed CS, which suggested that CS could alleviate stress in piglets.

Molecular oxygen is essential for the survival of all aerobic organisms. O₂ serves as the final electron acceptor in the process of aerobic energy metabolism, and reduces to H₂O by mitochondrial enzymes in the cell. However, partially reduced and highly reactive metabolites of O₂ may be formed when organisms suffer from endogenous and exogenous stimulation, to produce superoxide anion (O₂⁻) and hydrogen peroxide (H₂O₂). These derivatives of oxygen (reactive oxygen species, e.g. O₂⁻ and H₂O₂) are highly reactive and toxic, and can lead to increased oxidative stress in a variety of tissues. Oxidative stress can damage biomolecules, including cellular lipids, proteins, amino acids, and deoxyribonucleic acid, thereby inhibiting their normal function. For piglets, many endogenous and exogenous factors, such as environmental factors, weaning, biting, ranking, disease, and infection, can induce oxidative stress, resulting in poor growth performance.

Malondialdehyde is the most familiar breakdown product of lipid peroxidation, and its level is frequently used as a main marker to scale the extent of lipid oxidative damage caused by ROS. The authors observed that CS supplementation can decrease serum MDA content in weaned pigs, suggesting that CS could depress their lipid peroxidation. The scavenging capacity of free radicals is related to the non-enzymatic and enzymatic antioxidant defence systems, which were determined to further identify the manner of inhibition of oxidative damage by CS. Antioxidant enzymes are essential key factors against oxidative stress induced by

stimulus factors in animal defence system.

The enzyme defence system consists of SOD, CAT and GPx. SOD provides the efficient dismutation of superoxide radicals (O_2^-) into less toxic H_2O_2 , while CAT and GPx reduce H_2O_2 to O_2 and H_2O . Indeed, CS has shown beneficial effects on relieving oxidative stress induced by certain drugs or under various physiological and pathophysiological conditions. Ramasamy *et al.* reported that dietary supplementation with 200 mg CS/kg body weight per day for seven weeks, along with CCl_4 , markedly increased hepatic and circulatory SOD, CAT, GPx levels in rats. Additionally, Jeon *et al.* concluded that CS significantly increased hepatic CAT and SOD activities in chronic hepatic injuries in rats that had been induced by carbon tetrachloride. In addition, previous studies undertaken in the authors' laboratory showed that supplementation of CS in diets resulted in improvement in serum GPx, SOD and CAT activities in piglets and serum SOD activity in beef cattle. In agreement with those findings, the current results showed that dietary supplementation with CS increased serum SOD, GPx, and CAT activities, which are representative enzymatic antioxidants in the body.

The GSH is an important major non-enzymatic antioxidant in cells. Reduction of GSH could converse to oxidized GSH (GSSG) spontaneously or by enzymatic catalysis to trap radical species. In the current study, dietary CS increased GSH content in the serum of pigs, which was comparable with the study of Ramasamy *et al.*, which reported that dietary supplementation with CS resulted in higher GSH content in the livers of rats with hepatic injury induced by tetrachloride. Further, Anandan *et al.* reported that CS could have increased effects on the GSH level in the heart tissue of aged rats. This illustrates that CS as antioxidant could improve resistance to oxidation as measured by increased GSH content in serum.

The T-AOC, which includes a number of antioxidant enzymes and related biomolecules that can remove free radicals from a specific organ or living organism, reflects the total antioxidant ability. Limited published reports to date have shown that CS increased significantly the total antioxidant capacity of livestock. Obvious results in this study were observed in T-AOC level, while animals supplemented with CS also showed a dose-dependent effect. Thus, these results show that CS strengthened the oxidative defence mechanisms.

These combined findings suggest that CS might help to protect weaned pigs against oxidative stress, as they appear to reduce lipid peroxidation and improve their antioxidant defence systems functions *in vivo*.

With regard to the serum production of cytokines and immunoglobulins, which were used as further understanding of immune system status, the current study measured the concentrations of cytokines and immunoglobulins in serum. ILs are a group of cytokines that play an important role in the immune system. Among these, IL-1 β mediates the immune response by i) promoting the proliferation and differentiation of thymocytes and mature T-cells; ii) enhancing B-cell differentiation; iii) inhibiting the growth of tumour cells

and killing them; and (iv) inducing T-cells to generate IL-2. In addition, IL-2 is a key cytokine with broad spectrum crucial immunomodulatory activities. In the present study, CS was shown to increase the production of serum IL-1 β and IL-2 in piglets. Similar to the current findings, Yin *et al.* demonstrated that CS supplementation could promote the generation of IL-1 β of early-weaned piglet, since the increases in ILs indicate the activation of an immune reaction that might lead an improvement in immune status. Immunoglobulins are proteins with antibody activity that exist mainly in the plasma and are used to promote phagocytosis of monocytes and macrophages, and identify and neutralize bacteria and viruses. Serum IgG is the most persistent and important antibody in the primary immune response, and is used to promote phagocytosis of monocytes and macrophages, neutralize the toxicity of bacterial toxins, and combine with viral antigens to prevent the virus infecting host cells. Here, the authors reported that serum IgG level was increased markedly in CS supplemented pigs, indicating that CS efficiently modulated a humoral immune response, which may provide an advantage in increasing weaned pig immunity to prevent susceptibility to disease and enhance resistance to infections. The nitric oxide and arachidonic acid pathways were regulated by chitosan in a dose-dependent relationship may be reasons that chitosan affected the immune function in weaned piglets. All these results support the immunostimulatory effects of CS, which may be the reason for improved performance.

Conclusion

In conclusion, the results observed in this study showed that the use of a CS-supplemented diet may be effective in improving anti-oxidant and free radical scavenging capacity and regulating the immunological response of post-weaning piglets, and limiting the negative effects of weaning on growth performance in piglets.

References

- Calinescu, C., B. Mondovi, R. Federico, P. Ispas-Szabo and M.A. Mateescu. 2012. Carboxymethyl starch: Chitosan monolithic matrices containing diamine oxidase and catalase for intestinal delivery. *International Journal of Pharmaceutics*, 428 (1-2): 48-56.
- Han, B., X. Shi, Z. Gu, J. Xiao, X. Chu, J. Gu, Y. Zhang, J. Tan and Y. Ding. 2011. No enhanced expression and infectivity of porcine endogenous retrovirus in primary porcine hepatocytes with chitosan nanofiber scaffold. *Journal of Biomedical Nanotechnology*, 7 (3): 377-383.
- Henao-Diaz, Y.A., L. Gimenez-Lirola, K. Poonsuk, T. Cheng, C. Wang, J. Ji, D.H. Baum, R.G. Main and J.J. Zimmerman. 2018. Effect of chemical clarification of oral fluids on the detection of

- porcine reproductive and respiratory syndrome virus IgG. *Journal of Veterinary Diagnostic Investigation*, 30 (5): 671-677.
- Inaba, K., B.C. Branco, P. Rhee, B. Putty, O. Okoye, G. Barmparas, P. Talving and D. Demetriades. 2013. Long-term preclinical evaluation of the intracorporeal use of advanced local hemostatics in a damage-control swine model of grade IV liver injury. *Journal of Trauma and Acute Care Surgery*, 74 (2): 538-545.
- Lee, J.H., D.H. Lee, J.K. Park, S.K. Kim, C.H.D. Kwon and S.K. Lee. 2012. Effect of fulminant hepatic failure porcine plasma supplemented with essential components on encapsulated rat hepatocyte spheroids. *Transplantation Proceedings*, 44 (4): 1009-1011.
- Liu, G., S. Chen, G. Guan, J. Tan, N.A. Al-Dhabi, H. Wang, V. Duraipandiyar and J. Fang. 2016. Chitosan modulates inflammatory responses in rats infected with enterotoxigenic *Escherichia coli*. *Mediators of Inflammation*, 2016 (7432845).
- Liu, M., D. Yin, H. Fu, F. Deng, G. Peng, G. Shu, Z. Yuan, F. Shi, J. Lin, L. Zhao, L. Yin and G. Fan. 2017. Double-coated enrofloxacin microparticles with chitosan and alginate: Preparation, characterization and taste-masking effect study. *Carbohydrate Polymers*, 170: 247-253.
- Ly, G., L. Zhao, A. Zhang, W. Du, Y. Chen, C. Yu, X. Pan, Y. Zhang, T. Song, J. Xu, Y. Chen and L. Li. 2011. Bioartificial liver system based on choanoid fluidized bed bioreactor improve the survival time of fulminant hepatic failure pigs. *Biotechnology and Bioengineering*, 108 (9): 2229-2236.
- Siel, D., S. Vidal, R. Sevilla, R. Paredes, F. Carvallo, L. Lapiere, M. Maino, O. Perez and L. Saenz. 2016. Effectiveness of an immunocastration vaccine formulation to reduce the gonadal function in female and male mice by Th1/Th2 immune response. *Theriogenology*, 86 (6): 1589-1598.
- Xiong, Q., Y. Wei, H. Xie, Z. Feng, Y. Gan, C. Wang, M. Liu, F. Bai, F. Xie and G. Shao. 2014. Effect of different adjuvant formulations on the immunogenicity and protective effect of a live *Mycoplasma hyopneumoniae* vaccine after intramuscular inoculation. *Vaccine*, 32 (27): 3445-3451.
- Zhang, S., F. Zhang, B. Feng, Q. Fan, F. Yang, D. Shang, J. Sui and H. Zhao. 2015. Hematopoietic stem cell capture and directional differentiation into vascular endothelial cells for metal Stent-Coated Chitosan/Hyaluronic acid loading CD133 antibody. *Tissue Engineering Part a*, 21 (5-6): 1173-1183.